




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 307/87	A1	(11) International Publication Number: WO 00/39112 (43) International Publication Date: 6 July 2000 (06.07.00)
(21) International Application Number: PCT/DK99/00728 (22) International Filing Date: 22 December 1999 (22.12.99) (30) Priority Data: PA 1998 01718 23 December 1998 (23.12.98) DK (71) Applicant (for all designated States except US): H. LUNDBECK A/S [DK/DK]; Ottiliavej 9, DK-2500 Valby Copenhagen (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): PETERSEN, Hans [DK/DK]; Guldagervej 11, DK-2720 Vanløse (DK). DAHLBERG NIELSEN, Poul [DK/DK]; Prejlerupvej 26, DK-4560 Vig (DK). (74) Common Representative: H. LUNDBECK A/S; Petersen, John, Heidahl, Ottiliavej 9, DK-2500 Valby Copenhagen (DK).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: METHOD FOR THE PREPARATION OF 5-CYANOPHTHALIDE (57) Abstract <p>A method for the preparation of 5-cyanophthalide in which 5-carboxyphthalide is converted to the corresponding amide of Formula (IV) in which R is hydrogen or C₁₋₆ alkyl, which is then reacted with a dehydrating agent thereby obtaining 5-cyanophthalide. The conversion of 5-carboxyphthalide to the corresponding amide of Formula (IV) may be carried out via the corresponding C₁₋₆ alkyl or phenyl ester or the acid chloride, which is converted to the amide of Formula (IV) by amidation with ammonia or a C₁₋₆ alkylamine. By the process 5-Cyanophthalide, an important intermediate used in the preparation of the antidepressant drug citalopram, is prepared in high yields by a convenient, cost effective procedure.</p> <div style="text-align: center;">  </div> <div style="text-align: right;"> (IV) </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		


METHOD FOR THE PREPARATION OF 5-CYANOPHTHALIDE

The present invention relates to a novel process for the preparation of 5-cyanophthalide which is an intermediate used in the manufacture of the well known antidepressant drug


- 5 citalopram, 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofurancarbonitrile.

Background of the Invention.

- 10 Citalopram is a well known antidepressant drug that has now been on the market for some years and has the following structure:

**Formula I**

- 15 It is a selective, centrally active serotonin (5-hydroxytryptamine; 5-HT) reuptake inhibitor, accordingly having antidepressant activities. The antidepressant activity of the compound has been reported in several publications, eg. J. Hyttel, *Prog. Neuro-Psychopharmacol. & Biol. Psychiat.*, **1982**, 6, 277-295 and A. Gravem, *Acta Psychiatr. Scand.*, **1987**, 75, 478-486.
- 20 Citalopram is prepared by the process described in US Patent No 4,650,884, according to which 5-cyanophthalide is subjected to two successive Grignard reactions, *i.e.* with 4-fluorophenyl magnesium halogenide and N,N-dimethylaminopropyl magnesium halogenide, respectively, and the resulting compound of the formula

**Formula II**

is subjected to a ring closure reaction by dehydration with strong sulfuric acid.

Enantiomers of citalopram may be prepared by the method described in US Patent No. 4,943,590, i.e. by separating the enantiomers of the intermediate of Formula II and performing enantioselective ring closure in order to obtain the desired enantiomer.

- 5 Thus, 5-cyanophthalide is an important intermediate for the manufacture of citalopram and it is important to produce this material in an adequate quality, by a convenient process and in a cost-effective way.

- 10 A method for the preparation of 5-cyanophthalide has previously been described in Bull. Soc. Sci. Bretagne, 26, 1951, 35 and in Levy and Stephen, J. Chem. Soc., 1931, 867. By this method, 5-aminophthalide is converted to the corresponding 5-cyanophthalide by diazotation followed by reaction with CuCN. 5-Aminophthalide was obtained from 4-aminophthalimide by a two step reduction procedure.

- 15 Synthesis of certain alkyl- and phenylnitriles from acid chlorides is described in Tetrahedron Letters, 1982, 23, 14, 1505 - 1508, and in Tetrahedron, 1998, 54, 9281.

Though a number of other methods failed, it has been found that 5-cyanophthalide may be prepared in high yields by a convenient, cost-effective procedure from 5-carboxyphthalide.


20

Description of the invention

Accordingly, the present invention provides a novel method for the preparation of 5-cyanophthalide from 5-carboxyphthalide comprising

25


- a) converting 5-carboxyphthalide to an amide of Formula IV



in which R is hydrogen or C₁₋₆ alkyl, and


30

- b) then reacting the amide of Formula IV with a dehydrating agent thereby obtaining 5-cyanophthalide




Formula V

The conversion of 5-carboxyphthalide to the amide of Formula IV may be carried out via an ester of Formula VI or an acid chloride of Formula VII or via the ester and the acid chloride:



Formula VI



Formula VII

wherein R_1 is C_{1-6} alkyl or phenyl. The acid chloride is conveniently obtained by treatment of 5-carboxyphthalide with $POCl_3$, PCl_5 or $SOCl_2$ neat or in a suitable solvent, such as toluene or toluene comprising a catalytic amount of N,N -dimethylformamide. The ester is obtained by treatment of 5-carboxyphthalide with an alcohol R_1OH , wherein R_1 is as defined above, in the presence of an acid, preferably a mineral acid or a Lewis acid, such as HCl , H_2SO_4 , $POCl_3$, PCl_5 or $SOCl_2$. Alternatively, the ester may be obtained from the acid chloride by reaction with an alcohol. The ester of Formula VI or the acid chloride of Formula VII is then converted to the amide of Formula IV by amidation with ammonia or an C_{1-6} alkylamine, preferably t -butyl amine.

Throughout the specification and Claims, C_{1-6} alkyl refers to a branched or unbranched alkyl group having from one to six carbon atoms inclusive, such as methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl, 2,2-dimethyl-1-ethyl and 2-methyl-1-propyl.

The dehydrating agent used in step b) may be any suitable dehydrating agent, and the optimal agent may easily be determined by a person skilled in the art. Examples of suitable dehydrating agents are $SOCl_2$, $POCl_3$ and PCl_5 , preferably $SOCl_2$.

The reaction in step b) is carried out neat or in a suitable solvent, such as toluene, sulfolan or conveniently acetonitrile. When the reaction is carried out in a solvent, 1.0 - 1.5, preferably 1.0 - 1.2 equivalents of dehydrating agent is used per equivalent of the amide of Formula V. Furthermore, when a solvent is used, a catalytic amount of N,N -dimethylformamide may be needed, in particular when the dehydrating agent is $SOCl_2$. Preferably, toluene is used as the solvent, if necessary in the presence of a catalytic amount of N,N -dimethylformamide.

The reaction in step b) is carried out at elevated temperature, preferably at the reflux temperature of the solvent.

The reaction time is not important and may easily be determined by a person skilled in the art.

5

5-Cyanophthalide may be isolated in a conventional way, e.g. by addition of water, filtration and subsequent washing of the crystals. Further purification may, if desired, be performed by recrystallisation.

- 10 In a preferred embodiment of the process of the invention, R in Formula IV is H or t-butyl. When the reaction in step a) is carried out via an ester, R₁ is preferably methyl or ethyl.

- In a particularly preferred embodiment of the invention 5-carboxyphthalide of Formula III is reacted with an alcohol, R₁OH, preferably ethanol, in the presence of POCl₃, in order to obtain the corresponding ester of Formula VI, which is then reacted with ammonia thereby giving 5-carbamoylphthalide, which in turn is reacted with SOCl₂ in toluene comprising a catalytic amount of N,N-dimethylformamide.
- 15

- Surprisingly, substantially no reaction takes place at the lactone ring. Accordingly, by the process of the invention, 5-cyanophthalide is obtained in high yields and the process is much more convenient than the known process and uses more convenient and cheaper reactants and conditions.
- 20

- The 5-carboxyphthalide used as a starting material may be obtained by the methods described in US patent No. 3,607,884 or German patent No. 2630927, i.e. by reacting a concentrated solution of terephthalic acid with formaldehyde in liquid SO₃ or by electrochemical hydrogenation of trimellithic acid.
- 25

Examples

- 30 The invention is further illustrated by the following examples.

Example 1

Preparation of 5-Cyanophthalid

- 35 **5-Chlorocarbonylphthalid**

5-Carboxyphthalid (53 g, 0.3 mole) was suspended in toluene (200 mL) and thionylchloride (44 g, 0.6 mole). N,N-dimethylformamide (DMF) (1 mL) was added and the mixture was heated at reflux temperature for 3 hours. The mixture was cooled to room temperature and n-heptane was added (200 mL). The crystals formed were collected and washed with

heptane (100 mL). Yield 52 g, 88%. DSC onset: 131 °C. ¹H NMR (CDCl₃, 500 MHz): 5.47 (2H, s), 8.06 (1H, d, J = 7.5 Hz), 8.28 (1H, d, J = 7.5 Hz), 8.3 (1H, s). ¹³C NMR (CDCl₃, 125 MHz): 69.4, 125.1, 126.1, 131.1, 131.6, 137.8, 146.6, 167.4, 169.0.

5 **5-tert-Butylcarbamyolphthalid**

Method A):

5-Carboxyphthalid (36 g, 0.2 mole) was suspended in thionylchloride (100 mL). DMF (1.5 mL) was added and the mixture was refluxed for 1 hour. Toluene (200 mL) was added and the solvents were evaporated *in vacuo*. The residue was dissolved in tetrahydrofuran (THF) (200 mL) and added to a solution of tert-butylamine (31 g, 0.42 mole) in THF (200 mL) at 5 °C. The mixture was allowed to warm to room temperature and stirred overnight. The reaction was then poured into ice water (400 mL) and the precipitated crystals were filtered off. The crystals were washed with water (100 mL). Yield: 41 g, 87%. DSC onset: 189.5 °C.

15 Method B):

A solution of 5-chlorocarbonylphthalid (39 g, 0.2 mole) in THF (200 mL) was added to a solution of tert-butylamine (19 g, 0.25 mole) and triethylamine (26 g, 0.25 mole) in THF (200 mL) at room temperature. The mixture was stirred for 1 hour. The reaction mixture was then poured into ice water (500 mL). The crystalline material formed was collected and washed with water (100 mL).

Yield 42.5 g, 91%. DSC onset: 192 °C. Purity: 99.5% (hplc, peak area). ¹H NMR (DMSO-d₆, 500 MHz): 1.4 (9H, s), 5.46 (2H, s), 7.88 (1H, d, J = 7.5 Hz), 7.95 (1H, d, J = 7.5 Hz), 8.04 (1H, s). ¹³C NMR (DMSO-d₆, 125 MHz): 28.5, 51.2, 70.0, 122.0, 124.6, 126.6, 128.2, 141.3, 147.2, 165.5, 170.1.

25

5-Ethoxycarbonylphthalid

Method A):

5-Carboxyphthalid (37 g, 0.2 mole) was suspended in ethanol (400 mL). POCl₃ (10 g, 0.07 mole) was added drop-wise and the reaction mixture was heated to reflux temperature for 5 hours. Upon cooling to room temperature, the title compound crystallised. The crystals were filtered off and washed with ethanol (50 ml). Yield: 35 g, 87%. DSC onset: 151 °C. ¹H NMR (DMSO-d₆, 250 MHz): 1.36 (3H, t, J = 7 Hz), 4.38 (2H, q, J = 7 Hz), 5.48 (2H, s), 7.95 (1H, d, J = 7.5 Hz), 8.12 (1H, d, J = 7.5 Hz), . ¹³C NMR (DMSO-d₆, 62.5 MHz): 14.5, 61.5, 70.1, 124.0, 125.2, 128.8, 129.6, 134.8, 147.6, 164.9, 169.8.

35

Method B):

5-Chlorocarbonylphthalid (39 g, 0.2 mole) was suspended in ethanol (200 mL). The

mixture was heated to reflux for 15 minutes. After cooling, the crystalline material formed was filtered off and washed with ethanol (50 ml). Yield: 36 g, 88%. DSC onset: 151 °C.

5-Carbamylphthalid.

5 Method A):

5-Ethoxycarbonylphthalid (41 g, 0.2 mole) was suspended in ammonia (10M solution in methanol, 200 mL) in a pressure reactor. The reaction temperature was held at 80 °C for 20 hours. After cooling, the reaction mixture was poured onto ice (250 g) and pH was adjusted to pH=1 using concentrated hydrochloric acid. The mixture was stirred for 2 hours. The crystals formed were filtered off and washed with water (4x100 mL) and dried *in vacuo*. Yield: 33 g, 93%. DSC onset: 237 °C. ¹H NMR (DMSO-d₆, 250 MHz): 5.47 (2H, s), 7.65 (1H, s (NH)), 7.92 (1H, d, J = 7.5 Hz), 8.06 (1H, d, J=7.5 Hz), 8.14 (1H, s), 8.22 (1H, s (NH)). ¹³C NMR (DMSO-d₆, 62.5 MHz): 70.0, 122.2, 124.9, 127.2, 128.2, 139.7, 147.4, 167.1, 170.1.

15

Method B):

5-Chlorocarbonylphthalid (20 g, 0.1 mole) was dissolved in THF (100 mL) and added to ammonium hydroxide (50 mL) in ice water (300 mL). The mixture was stirred for 30 minutes and the precipitated crystals were filtered off. The crystals were washed with water (100 mL) and dried *in vacuo*. Yield: 17.1 g, 97%. DSC onset: 237 °C.

20

5-Cyanophthalid.

Method A):

Dry 5-carbamylphthalid (36 g, 0.2 mole) was suspended in toluene (600 mL) and thionylchloride (36 g, 0.3 mole) was added. DMF (2 mL) was added. The reaction mixture was heated at 75 °C for 6 hours. Toluene (100 mL) was removed by distillation and the remaining solution was cooled to room temperature. The crystals formed were filtered off and washed with toluene (150 mL) and water (100 mL). The product was recrystallised from toluene. Yield: 22 g, 80%. DSC onset: 203 °C.

30

Method B):

Tert.-Butylcarbamylphthalid (23.3 g, 0.1 mole) was suspended in thionylchloride (100 mL). The mixture was heated to reflux for 30 min. Toluene (100 mL) was added and the solvents were removed *in vacuo*. The title product was crystallised from acetic acid or toluene.


35

Yield 15.5 g, 93% from toluene. DSC onset: 203 °C. Purity: 98% (hplc, peak area).

CLAIMS


1. A method for the preparation of 5-cyanophthalide comprising

- 5 a) conversion of 5-carboxyphthalide to an amide of Formula IV




in which R is hydrogen or C₁₋₆ alkyl, and

- 10 b) then reacting the amide of Formula IV with a dehydrating agent thereby obtaining 5-cyanophthalide




- 15 2. The method of Claim 1 wherein the conversion of 5-carboxyphthalide to the amide of Formula IV is carried out via an ester of Formula VI;



- 20 wherein R₁ is C₁₋₆ alkyl or phenyl, by treatment of 5-carboxyphthalide with an alcohol R₁OH in the presence of an acid and subsequent amidation of the ester of formula VI with ammonia or an C₁₋₆ alkylamine.

3. The method of Claim 1 wherein the conversion of 5-carboxyphthalide to the amide of Formula IV is carried out via an acid chloride of Formula VII:

8




Formula VII


by treatment of 5-carboxyphthalide with POCl_3 , PCl_5 or SOCl_2 and subsequent amidation of the acid chloride of formula VII with ammonia or an C_{1-6} alkylamine.

5

4. The method of Claim 1 wherein the conversion of 5-carboxyphthalide to the amide of Formula IV is carried out via an acid chloride of Formula VII and an ester of Formula VI:



Formula VI



Formula VII

10 wherein R_1 is C_{1-6} alkyl or phenyl, by treatment of 5-carboxyphthalide with POCl_3 , PCl_5 or SOCl_2 , reacting the acid chloride of Formula VII thus formed with an alcohol R_1OH and performing amidation of the ester of Formula VI with ammonia or an C_{1-6} alkylamine.

15 5. The method of Claim 2 wherein the acid used is a mineral acid or a Lewis acid, preferably HCl , H_2SO_4 , POCl_3 , PCl_5 , SOCl_2 .

6. The method of Claim 2, 4 or 5 wherein R_1 is methyl or ethyl.

20 7. The method of any of Claims 1 - 6 in which the dehydrating agent used in step b) is SOCl_2 , POCl_3 or PCl_5 , preferably SOCl_2 .

8. The method of any of Claims 1 - 7 wherein the reaction in step b) is carried out neat or in a suitable solvent, such as toluene, sulfolan or acetonitrile, preferably in toluene.

25 9. The method of any of Claim 8 wherein the dehydrating agent used in step b) is SOCl_2 and the reaction is carried out in toluene comprising a catalytic amount of N,N -dimethylformamide.

30 10. The method of any of Claims 1-9 wherein R is H or tert-butyl

11. The method of Claim 2, wherein the 5-carboxyphthalide of Formula III is reacted with an alcohol R_1OH , preferably ethanol or methanol, in the presence of $POCl_3$, in order to obtain an ester of Formula VI, which is then reacted with ammonia, thereby giving 5-carbamoylphthalide, which in turn is reacted with $SOCl_2$ to 5-cyanophthalide.

5

12. The method of Claim 11, wherein the 5-carboxyphthalide of Formula III is reacted with ethanol in the presence of $POCl_3$, in order to obtain the ethyl ester of Formula VI, which is then reacted with ammonia in methanol, thereby giving 5-carbamoylphthalide, which in turn is reacted with $SOCl_2$ to 5-cyanophthalide.

10

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00728

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 307/87

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0171943 A1 (H. LUNDBECK A/S), 19 February 1986 (19.02.86), formula III --	1-12
Y	WO 9819513 A2 (H. LUNDBECK A/S), 14 May 1998 (14.05.98), formula IV --	1-12
Y	Calvin A. Buehler et al, "Survey of Organic syntheses", Wiley-Interscience, A division of John Wiley & Sons, Inc., page 951 --	1-12

☒ Further documents are listed in the continuation of Box C.
 ☒ See patent family annex.

* Special categories of cited documents

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

31 March 2000

Date of mailing of the international search report

03 -05- 2000

Name and mailing address of the ISA

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Göran Karlsson/Eö

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00728

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>Sir Derek Barton, F.R.S. et al, "Comprehensive Organic Chemistry. The Synthesis and Reactions of Organic Compounds", Volume 2, Edited by I. O. Sutherland, pages 1024-1025</p> <p>-- -----</p>	1-12

INTERNATIONAL SEARCH REPORT

Information on patent family members

02/12/99

International application No.

PCT/DK 99/00728

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0171943 A1	19/02/86	SE 0171943 T3	
		AT 38661 T	15/12/88
		AU 574819 B	14/07/88
		AU 4577685 A	13/02/86
		CA 1237147 A	24/05/88
		DE 3566251 A	22/12/88
		DK 89595 A	10/08/95
		DK 356285 A	07/02/86
		ES 545885 A	01/04/86
		FI 81338 B,C	29/06/90
		FI 852902 A	07/02/86
		GR 851894 A	03/12/85
		IE 57817 B	21/04/93
		IL 75690 A	31/10/88
		JP 1902596 C	08/02/95
		JP 6025099 B	06/04/94
		JP 61087654 A	06/05/86
		NO 853091 A	07/02/86
		NZ 212541 A	30/05/88
		PT 80913 A,B	01/09/85
		US 4650884 A	17/03/87
WO 9819513 A2	14/05/98	AU 6609898 A	29/05/98